

AcrySof® IQ ReSTOR® +3.0 D Multipiece Multifocal IOL



MN6AD1

- AcrySof® IQ ReSTOR® +3.0 D Multifocal IOL is available in a multipiece option (MN6AD1)
 - Suggested A-constant is 119.2 for capsular bag placement
 - Multipiece design
 - Available in 6.0-30.0 D in half diopter
 - Available in 31.0-34.0 D in whole diopter
 - Apodized diffractive optic identical to the leading single piece design of the SN6AD1 model
- Currently qualified for B-cartridge and MONARCH® II injector for implantation



AcrySof® IQ ReSTOR® +3.0 D Multifocal IOLs

Specifications for AcrySof® IQ ReSTOR® IOL Models

Model number	SN6AD1	MN6AD1
Design	single piece	multi-piece
Add power	+3.0 D	+3.0 D
Add power to spectacle plane	+2.5 D	+2.5 D
Number of steps	9	9
Diopter range	+6.0 to +30.0 D +31.0 to +34.0 D (1 diopter increments)	+6.0 to +30.0 D +31.0 to +34.0 D (1 diopter increments)
Optic type	Apodized, Diffractive, Aspheric	Apodized, Diffractive, Aspheric
Optic diameter	6.0 mm	6.0 mm
Overall length	13.0 mm	13.0 mm
Haptic angulation	0°	10°
Suggested A-constant	118.9	119.2
Filtration	UV and Blue-Light Filtering	UV and Blue-Light Filtering



MN6AD1



SN6AD1

AcrySof® IQ ReSTOR® IOL IMPORTANT SAFETY INFORMATION

CAUTION: Federal (USA) law restricts this device to the sale by or on the order of a physician.

INDICATIONS: The AcrySof® IQ ReSTOR® Posterior Chamber Intraocular Lens (IOL) is intended for primary implantation for the visual correction of aphakia secondary to removal of a cataractous lens in adult patients with and without presbyopia, who desire near, intermediate and distance vision with increased spectacle independence. The lens is intended to be placed in the capsular bag.

WARNING/PRECAUTION: Careful preoperative evaluation and sound clinical judgment should be used by the surgeon to decide the risk/benefit ratio before implanting a lens in a patient with any of the conditions described in the Directions for Use labeling. Physicians should target emmetropia, and ensure that IOL centration is achieved. Care should be taken to remove viscoelastic from the eye at the close of surgery.

Some patients may experience visual disturbances and/or discomfort due to multifocality, especially under dim light conditions. Clinical studies with the AcrySof® ReSTOR® lens indicated that posterior capsule opacification (PCO), when present, developed earlier into clinically significant PCO. Prior to surgery, physicians should provide prospective patients with a copy of the Patient Information Brochure available from Alcon for this product informing them of possible risks and benefits associated with the AcrySof® IQ ReSTOR® IOLs.

Studies have shown that color vision discrimination is not adversely affected in individuals with the AcrySof® Natural IOL and normal color vision. The effect on vision of the AcrySof® Natural IOL in subjects with hereditary color vision defects and acquired color vision defects secondary to ocular disease (e.g., glaucoma, diabetic retinopathy, chronic uveitis, and other retinal or optic nerve diseases) has not been studied. Do not resterilize; do not store over 45° C; use only sterile irrigating solutions such as BSS® or BSS PLUS® Sterile Intraocular Irrigating Solutions.

ATTENTION: Reference the Directions for Use labeling for a complete listing of indications, warnings and precautions.

Alcon®

a Novartis company

Contact your Alcon surgical representative for more information

© 2013 Novartis 1/13 RES13017SS